



(REVIEW ARTICLE)



NEUROPSYCHOLOGICAL ANALYSIS OF THE CASE OF DR. P. IN OLIVER SACK'S "THE MAN WHO MISTOOK HIS WIFE FOR A HAT"

Penny Lagonika *

Net Media Lab, Institute of Informatics and Telecommunications, NCSR Demokritos, Athens, Greece.

* Corresponding Author

Global Journal of Engineering and Technology Advances, 2026, 28(03), 256–262

Publication history: Received on 10 August 2026; revised on 19 September 2026; accepted on 21 September 2026

Article DOI: <https://doi.org/10.30574/gjeta.2026.28.3.0259>

Copyright © 2026 Author(s) retain the copyright of this article. This article is published under the terms of the Creative Commons Attribution License 4.0

ABSTRACT

The present paper addresses disorders of visual perception and recognition through the analysis of the classic neuropsychological case of Dr. P., as described by Oliver Sacks. Although the patient retains normal vision, intelligence, and language abilities, he exhibits a severe inability to recognize faces and objects, as well as to assign meaning to visual stimuli. The study focuses on the distinction between the forms of visual agnosia, as formulated by Lissauer, and highlights the importance of integrative processing for the holistic perception of the visual world. Particular emphasis is placed on the role of neuroplasticity and compensatory mechanisms, with music functioning as a primary means of organizing the patient's experience and adapting to everyday life.

Keywords: Visual Agnosia, Integrative Agnosia, Prosopagnosia, Ventral Visual Stream, Neuroplasticity, Oliver Sacks, Neuropsychology

1 INTRODUCTION

Neuropsychology is a branch of psychology that studies the relationship between the structure and function of the brain and behavior, cognition, and emotion. Through the study of patients with focal or degenerative brain damage, neuropsychology provides valuable insights into the organization of higher cognitive functions. One of the most famous cases in the neuropsychological literature is that of Dr. P., as described by Oliver Sacks, which concerns a severe disorder of visual perception known as visual agnosia.

The purpose of this paper is to present an analytical account of this case study, describe the disorder exhibited by the patient, investigate the dysfunctional neural substrates, and refer to therapeutic and rehabilitation approaches according to the theoretical framework of contemporary neuropsychology.

2 CASE STUDY AND NEUROLOGICAL ASSESSMENT

Dr. P. is the patient being examined and presents a particularly complex disorder of visual perception. He is an educated, socially functional, and emotionally stable individual, with no indications of generalized dementia or cognitive decline. His difficulties are exclusively related to the way in which he perceives and interprets the visual world, whereas his memory and thought process appears to remain unaffected.

The first concerning signs appear in his everyday life, when he fails to recognize faces, even familiar ones, such as his students or his wife. He does not recognize faces as unified wholes but instead focuses fragmentarily on individual features. He does not understand facial expressions, nor can he assign emotional meaning to them, which constitutes a clear indication of prosopagnosia. He frequently confuses inanimate objects with living things, such as when he attempts to put his wife's head on, believing it to be a hat.

Initially, Dr. P. consults an ophthalmologist because he is concerned about possible visual complications related to his diagnosis of diabetes. However, after a detailed examination, the doctor determines that his eyes are healthy and concludes that the problem is not ophthalmological but is located in the visual areas of the brain. This observation is crucial, as it confirms that the disorder involves higher-level processing of visual information. During the neurological examination, a mild abnormality in reflexes is observed on his left side, as it becomes apparent that the patient has difficulty perceiving stimuli located in his left visual field. In addition, he is unable to perceive images in their entirety, he perceives only individual details without integrating them into a unified, meaningful whole. Nothing appears visually familiar to him. Objects and faces are approached in a mechanical manner, as though they were abstract problems to be analyzed. To recognize objects, Dr. P. often relies on other senses, mainly smell and hearing. A rose, for example, is recognized only when he smells it. He also appears unable to recall visual characteristics from memory, although he has no difficulty understanding situations, plots, and dialogues. He remembers events and narratives, but without any visual representation. The pattern clearly suggests a form visual agnosia. Even his dreams lack visual content, meaning is conveyed without images, through non-visual sensory or conceptual forms.

3 VISUAL AGNOSIA: DEFINITION AND CATEGORIES

The term visual agnosia is described within the theoretical framework initially established by Lissauer, whose work laid the foundation for the modern understanding of the condition.. Heinrich Lissauer was a German neurologist, born in 1861 and active in the late nineteenth century. Specifically, in his article *Ein Fall von Seelenblindheit* (1890), he

analyzed in detail the case of a patient who, similarly to Dr. P. in Oliver Sacks's account, was unable to visually recognize even common objects, while being able to recognize them through touch or hearing. He initially referred to this condition as *Seelenblindheit* (meaning "mind blindness"). The condition is defined as a state in which a patient, although receiving visual impressions with sufficient acuity to distinguish stimuli, is unable to correctly recognize and name objects through vision or provide information about their properties (Lissauer, 1980).

According to Lissauer, the classification of visual agnosia is based on the analysis of the recognition process into two distinct mental stages: perception and association. Perception is the act of consciously becoming aware of a sensory impression. At this stage, the individual perceives the form and details of the object as sensory data. Association concerns the connection of the content of perception with other pre-existing representations and concepts in memory. Through this process, the object acquires meaning (Serino, 2014).

In this way, two forms of the disorder emerge: Associative Visual Agnosia and Apperceptive Visual Agnosia. A useful analogy for understanding this distinction becomes apparent through the act of reading. In visual agnosia, although the language is known, the letters appear as individual lines that never form words, whereas in associative agnosia, the image of the word is clear but its meaning has been forgotten, even though the concept is known when the word is heard.

In the case of Dr. P., the ophthalmologist concluded that his eyes were functioning perfectly, but his brain was unable to interpret the images. This is consistent with research indicating that visual acuity and visual fields may be normal, while the impairment is located at higher levels of processing in the brain (Riddoch & Humphreys, 2003). It is important, however, to note that despite the distinction between the two disorders, in clinical practice the two forms are rarely completely separate. Lissauer himself argues that purely associative agnosia is almost contradictory, since damage to associative processes inevitably also affects the clarity of perception, particularly in complex forms.

Regarding Dr. P.'s difficulty in perceiving objects located in his left visual field, it is noted that there is an important clinical association between visual agnosia, particularly prosopagnosia, and left homonymous hemianopia (Meadows, 1974).

4 INTEGRATIVE AGNOSIA AND PROSOPAGNOSIA

Integrative agnosia is a form of visual agnosia characterized by the patient's inability to combine the individual elements of a visual stimulus into a single, coherent form (Riddoch & Humphreys, 1987). One of the most characteristic findings described is the phenomenon of over-segmentation. Patients perceive the individual details of an object as independent elements, without being able to integrate them. Thus, an entire object with distinct functionality may be described as a

collection of unrelated parts. This inability is closely associated with dysfunction in the process of “grouping,” as described by Behrmann & Williams (2007). In integrative agnosia, patients can process only a very limited number of individual elements simultaneously and experience considerable difficulty with complex stimuli.

Similarly, prosopagnosia is defined as the inability to recognize familiar people despite the preservation of general intelligence and visual acuity. This disorder may occur either as an isolated phenomenon or as part of a broader visual agnosia (Kay & Levin, 1982). Prosopagnosia is classically divided into two main forms: acquired and congenital or developmental. Acquired prosopagnosia occurs following brain damage, such as a stroke, traumatic injury, anoxia, or neurodegenerative disease, in individuals who previously possessed normal face-recognition abilities. The lesions are often located in the temporal and occipital cortex, particularly in areas involved in processing the visual identity of faces (Barton, 2008).

In contrast, congenital or developmental prosopagnosia is a lifelong neurodevelopmental disorder in which the individual fails to develop effective face-recognition abilities without a history of obvious brain damage. These individuals have normal vision, intelligence, and memory, but display a selective impairment in face recognition from childhood. Furthermore, they often lack awareness of this impairment, as they have developed compensatory strategies, such as recognition through voice or movement, and consider their way of perceiving people to be normal (Grüter et al., 2008; Behrmann & Avidan, 2005).

5 DR. P.'S PROSOPAGNOSIA

In Dr. P.'s case, there appears to be a severe disorder of visual recognition that extends beyond the boundaries of isolated prosopagnosia. Although Dr. P. has a difficulty recognizing his students when they approach him, which could initially suggest congenital prosopagnosia, the clinical picture indicates a broader and deeper deficit. His deficit is not limited to recognizing faces but also extends to recognizing inanimate objects, which he often confuses with human features, such as when he famously mistook his wife's head for a hat (Sacks, 1985). This is described as an indication of a severe disorder in the processing of visual form rather than an isolated inability to recognize faces (Behrmann & Williams, 2007). In this sense, Dr. P.'s prosopagnosia does not appear to constitute an independent symptom, but rather to arise secondarily from the generalized associative visual agnosia that he presents. The patient is unable to integrate individual visual features into a coherent and semantically meaningful whole, resulting in objects and faces lacking recognizable identity for him. As Lissauer (1980) had already pointed out, for the agnostic patient, facial expressions cease to have symbolic meaning and become a “combination of incomprehensible arabesques.” As a compensatory strategy, the patient is forced to examine individual visual elements in a fragmented manner, as becomes evident when Dr. P. is unable to recognize a glove and describes it as “some kind of case.”

This interpretation is further supported by Dr. P.'s difficulty in recognizing human emotions through facial expressions. According to the neuropsychological literature, emotion recognition and face recognition are distinct - although interconnected - functions supported by different neural networks (Calder & Young, 2005). However, in cases of generalized agnosia, the failure of holistic form processing affects both abilities. It is also reported, that in 80.3% of cases, deficits in face and object recognition occur together (Behrmann & Geskin, 2017). This data supports the notion that, at least in some patients, prosopagnosia is not an isolated disorder but rather part of a broader integrative agnosia.

6 NEUROLOGICAL BASIS

The neurological basis of visual agnosia, and particularly of the integrative form, is located mainly in the posterior regions of the brain, which participate in the successive stages of visual processing. Extensive lesions in the occipital cortex, which includes the primary and secondary visual cortex (Brodmann areas 17, 18, and 19), can disrupt basic mechanisms involved in the analysis of visual stimuli, such as sensitivity to line orientation and edge detection, functions that are necessary for constructing the form of objects (Hubel & Wiesel, 1968).

Visual agnosia has been particularly associated with lesions in the ventral visual stream, also known as the “what” system, which extends from the occipital to the temporal cortex and is responsible for the recognition and semantic processing of objects (Goodale & Milner, 1992). Since Lissauer, it has been proposed that agnosia may result not only from focal cortical lesions but also from disruption of the connections between visual areas and higher associative cortical networks, leading to a failure to integrate visual information (Lissauer, 1890). Integrative agnosia, in particular, is associated with lesions in the inferior occipito-temporal cortex, an area that is critical for binding and combining individual visual features into unified representations. Studies of patients have shown bilateral lesions in the fusiform and lingual gyri, regions that appear to play a central role in integrating form and overcoming fragmented processing (Behrmann & Kimchi, 2003).

Lastly, prosopagnosia occurring as part of a broader disorder, has been mainly associated with lesions in the posterior portion of the right temporal lobe and, more specifically, in the fusiform gyrus. This region is considered crucial for the specialized processing of faces, and its dysfunction leads to an inability to recognize faces holistically despite the preservation of individual visual information (Barton, 2008). Specifically, in the case of Dr. P., Sacks reports a damaged cerebral cortex as a result of a possible large brain tumor or a degenerative process in the visual areas of the brain.

7 NEUROPLASTICITY: MUSIC AS A COMPENSATORY MECHANISM IN DR. P.'S EVERYDAY LIFE

In the case of Dr. P., while his visual processing shows severe dysfunction due to damage to the ventral visual stream and the occipito-temporal cortex, his auditory and musical networks remain not only intact, but also highly developed. Music is not merely a simple interest for Dr. P.; rather, it permeates his entire daily life and functions as a central organizing principle of his experience and behavior. Neuroanatomically, musical ability and perception are associated with a broad network of brain regions that includes auditory areas as well as systems involved in memory, emotion, movement, and reward. The recognition of music and tones is supported by a broad network that includes the superior temporal gyrus (BA 22/42) in both hemispheres for initial auditory analysis (Zatorre et al., 1998). The processing of melody, rhythm, and pitch also activates regions of the temporal lobe, prefrontal cortex, cerebellum, and basal ganglia, networks associated with memory, prediction, and movement (Shulman et al., 1993). Structures related to emotion, such as the hippocampus and the cingulate cortex, also participate, helping explain why music can evoke strong emotions (Schlaug et al., 2006).

The temporal, frontal, and subcortical structures involved in musical processing do not overlap completely with the visual systems responsible for recognizing objects and faces. Thus, while Dr. P. is unable to integrate visual stimuli into coherent forms, he can accurately process rhythm, melody, and harmony. Furthermore, available evidence in literature suggests that individuals with focal brain dysfunctions often exhibit compensation through increased functioning of other networks, a phenomenon attributed to neuroplasticity and cross-regional reorganization of the brain (Chen et al., 2010). For example, musical abilities in people with cognitive impairments or Alzheimer's disease may remain intact or even become enhanced due to the preservation or strengthening of other brain networks (Matziorinis et al., 2024). In Dr. P.'s case, music appears to assume this compensatory role.

Within this context frame, Sacks's observation that "music had taken the place of image" is particularly significant. This phrase aptly summarizes the functional substitution that appears to have occurred: where vision fails to provide recognizable forms, music assumes the role of organizing experience and connecting the individual with the world. For Dr. P., reality is no longer structured primarily visually, but rather through sound and rhythm. Thus, his strong musical inclination is not merely a parallel characteristic of his personality, but a central element of adaptation and survival within the visual agnosia he experiences.

8 THERAPEUTIC APPROACHES AND REHABILITATION METHODS

Although there is no established, universally effective treatment for all types of visual agnosia, there are valid therapeutic and rehabilitation strategies that can improve functioning and quality of life in individuals with these disorders. These approaches focus primarily on strengthening residual functioning and compensatory mechanisms rather than complete neural restoration (Heutink et al., 2019). More specifically, these compensatory mechanisms include training in alternative methods of object recognition, adapting the environment to facilitate functioning, and using other sensory information, such as touch or sound, to compensate for visual dysfunction. For example, a case study showed that a patient with integrative visual agnosia could improve independence in daily activities through training that utilized preserved tactile abilities, such as touch, and repeated learning, although generalization of the skills outside the living environment required continuous support (Tanaka, 2021).

Other contemporary therapeutic procedures include the use and exploitation of digital technologies, which are used extensively in education and rehabilitation domains. Therefore, the importance of all digital technologies in the field of education is highlighted in our final point. ICTs support universal access to education, provide innovative approaches for effective teacher training, enhance learning retention, promote cooperation, increase openness, develop learner-centered approaches, and hasten the process of learning. Additionally, by using virtualization, mobilization, artificial intelligence, and new learning environments like virtual worlds, support educational activities and methodologies. More specifically, ICTs are very effective and productive in psychological training, facilitating and improving the assessment, intervention, and educational procedures via mobile devices that bring educational activities everywhere [24] and through a variety of ICT applications that serve as the backbone of education [25-28]. While games turn education into a multimodal, incredibly amiable, and enjoyable interaction, the use of AI, STEM, and ROBOTICS raise educational procedures into new levers of adaptation, creativity, and performance [29-34]. Furthermore, the adoption, improvement, and fusion of ICTs with theories and models of metacognition, mindfulness, meditation, and emotional intelligence cultivation [35-37] places the development of mental abilities at the center of educational procedures and

policies, which accelerates and improves educational practices and outcomes, particularly in psychological rehabilitation procedures.

9 CONCLUSIONS

The case of Dr. P. represents a characteristic example of generalized visual agnosia with prominent features of integrative agnosia and secondary prosopagnosia. Through a detailed analysis of the patient's symptoms, it becomes clear that the deficit does not concern visual acuity or general cognitive functioning, but rather the higher-level processing and integration of visual information, which significantly affects his everyday functioning and interaction with the world. At the same time, the analysis highlighted the anatomical pathophysiology of the disorder, as well as the importance of neuroplasticity. Music functions for Dr. P. as a primary compensatory mechanism, organizing his experience and facilitating his adaptation to everyday life. Overall, the case highlights the complexity of visual agnosia and the need for a holistic approach to diagnosis and rehabilitation. Sacks himself notes how fascinated he was by the uniqueness of Dr. P., a man who had almost completely lost the world as a visual representation, but had preserved it intact as music and as will. Despite the gradual progression of the disease, Dr. P. continued to live, teach, and give meaning to his existence through music. His story is a remarkable example of human psychological as well as biological resilience.

STATEMENTS ON COMPLIANCE WITH ETHICAL STANDARDS

Acknowledgments

The Authors would like to thank the SPECIALIZATION IN ICTs AND SPECIAL EDUCATION: PSYCHOPEDAGOGY OF INCLUSION Postgraduate studies Team, for their support.

Disclosure of conflict of interest

The Authors proclaim no conflict of interest.

REFERENCES

- [1] Barton, J. J. S. (2008). Structure and function in acquired prosopagnosia: Lessons from a series of 10 patients with brain damage. *Journal of Neuropsychology*, 2(1), 197–225. <https://doi.org/10.1348/174866407X214172>
- [2] Behrmann, M., & Avidan, G. (2005). Congenital prosopagnosia: Face-blind from birth. *Trends in Cognitive Sciences*, 9(4), 180–187. <https://doi.org/10.1016/j.tics.2005.02.011>
- [3] Behrmann, M., & Kimchi, R. (2003). What does visual agnosia tell us about perceptual organization and its relationship to object perception? *Journal of Experimental Psychology: Human Perception and Performance*, 29(1), 19–42. <https://doi.org/10.1037/0096-1523.29.1.19>
- [4] Behrmann, M., & Williams, P. (2007). Impairments in part-whole representations of objects in two cases of integrative visual agnosia. *Cognitive Neuropsychology*, 24(7), 701–730. <https://doi.org/10.1080/02643290701672764>
- [5] Calder, A. J., & Young, A. W. (2005). Understanding the recognition of facial identity and facial expression. *Nature Reviews Neuroscience*, 6(8), 641–651. <https://doi.org/10.1038/nrn1724>
- [6] Chen, H., Epstein, J., & Stern, E. (2010). Neural plasticity after acquired brain injury: Evidence from functional neuroimaging. *PM & R: The Journal of Injury, Function, and Rehabilitation*, 2(12 Suppl 2), S306–S312. <https://doi.org/10.1016/j.pmrj.2010.10.006>
- [7] Geskin, J., & Behrmann, M. (2017). Congenital prosopagnosia without object agnosia? A literature review. *Cognitive Neuropsychology*. <https://doi.org/10.1080/02643294.2017.1392295>
- [8] Goodale, M. A., & Milner, A. D. (1992). Separate visual pathways for perception and action. *Trends in Neurosciences*, 15(1), 20–25. [https://doi.org/10.1016/0166-2236\(92\)90344-](https://doi.org/10.1016/0166-2236(92)90344-)
- [9] Grüter, T., Grüter, M., & Carbon, C. C. (2008). Neural and genetic foundations of face recognition and prosopagnosia. *Journal of Neuropsychology*, 2(1), 79–97. <https://doi.org/10.1348/174866407X231001>
- [10] Heutink, J., Indorf, D. L., & Cordes, C. (2019). The neuropsychological rehabilitation of visual agnosia and Balint's syndrome. *Neuropsychological Rehabilitation*, 29(10), 1489–1508. <https://doi.org/10.1080/09602011.2017.1422272>

- [11] Hubel, D. H., & Wiesel, T. N. (1968). Receptive fields and functional architecture of monkey striate cortex. *The Journal of Physiology*, 195(1), 215–243. <https://doi.org/10.1113/jphysiol.1968.sp008455>
- [12] Kay, M. C., & Levin, H. S. (1982). Prosopagnosia. *American Journal of Ophthalmology*, 94(1), 75–80. [https://doi.org/10.1016/0002-9394\(82\)90194-5](https://doi.org/10.1016/0002-9394(82)90194-5)
- [13] Lissauer, H. (1890). Ein Fall von Seelenblindheit nebst einem Beitrage zur Theorie derselben. [A case of mindblindness and a contribution to the theory]. *Archiv für Psychiatrie und Nervenkrankheiten*, 21(1), 222–270. <https://doi.org/10.1007/BF02226765>
- [14] Matziorinis, A. M., Leemans, A., Skouras, S., Flo, B. K., Bashevkin, T., & Koelsch, S. (2024). The effects of musicality on brain network topology in the context of Alzheimer’s disease and memory decline. *Imaging Neuroscience*. https://doi.org/10.1162/imag_a_00248
- [15] Meadows, J. C. (1974). Disturbed perception of colours associated with localized cerebral lesions. *Brain*, 97(1), 615–632. <https://doi.org/10.1093/brain/97.1.615>
- [16] Riddoch, M. J., & Humphreys, G. W. (1987). Visual object processing in optic aphasia: A case of semantic access agnosia. *Cognitive Neuropsychology*, 4(2), 131–185. <https://doi.org/10.1080/02643298708252038>
- [17] Riddoch, M. J., & Humphreys, G. W. (2003). Visual agnosia. *Neurologic Clinics*, 21(2), 501–520. [https://doi.org/10.1016/S0733-8619\(02\)00095-6](https://doi.org/10.1016/S0733-8619(02)00095-6)
- [18] Sacks, O. (1985). *The Man Who Mistook His Wife for a Hat*. Harper & Row.
- [19] Schlaug, G., Marchina, S., & Norton, A. (2009). Evidence for plasticity in white-matter tracts of patients with chronic Broca’s aphasia undergoing intense intonation-based speech therapy. *Annals of the New York Academy of Sciences*, 1169(1), 385–394. <https://doi.org/10.1111/j.1749-6632.2009.04587.x>
- [20] Serino, A., Cecere, R., Dundon, N., Bertini, C., Sanchez-Castaneda, C., & Làdavas, E. (2014). When apperceptive agnosia is explained by a deficit of primary visual processing. *Cortex*, 52, 12–27. <https://doi.org/10.1016/j.cortex.2013.12.011>
- [21] Shulman, R. G., Blamire, A. M., Rothman, D. L., & McCarthy, G. (1993). Nuclear magnetic resonance imaging and spectroscopy of human brain function. *Proceedings of the National Academy of Sciences of the United States of America*, 90(8), 3127–3133. <https://doi.org/10.1073/pnas.90.8.3127>
- [22] Tanaka, M. (2021). Occupational therapy in integrative visual agnosia and memory disorder: A case report. *Asian Journal of Occupational Therapy*, 17(1), 27–30. <https://doi.org/10.11596/asijot.17.27>
- [23] Zatorre, R. J., Perry, D. W., Beckett, C. A., Westbury, C. F., & Evans, A. C. (1998). Functional anatomy of musical processing in listeners with absolute pitch and relative pitch. *Proceedings of the National Academy of Sciences*, 95(6), 3172–3177. <https://doi.org/10.1073/pnas.95.6.3172>
- [24] A Drigas, DE Dede, S Dedes 2020 Mobile and other applications for mental imagery to improve learning disabilities and mental health *International Journal of Computer Science Issues (IJCSI)* 17 (4), 18-23
- [25] Pappas, M.A., & Drigas, A.S. (2015). ICT based screening tools and etiology of dyscalculia. *International Journal of Engineering Pedagogy*, 3, 61-66.
- [26] Drigas, A., & Kostas, I. (2014). On Line and other ICTs Applications for teaching math in Special Education. *International Journal of Recent Contributions from Engineering, Science & IT (iJES)*, 2(4), pp-46. <http://dx.doi.org/10.3991/ijes.v2i4.4204>
- [27] Lizeta N. Bakola, Nikolaos D. Rizos, Drigas, A. S., “ICTs for Emotional and Social Skills Development for Children with ADHD and ASD Co-existence” *International Journal of Emerging Technologies in Learning (iJET)*, <https://doi.org/10.3991/ijet.v14i05.9430>
- [28] Drigas, A. S., Koukianakis, L, Papagerasimou, Y. (2006) “An elearning environment for nontraditional students with sight disabilities.”, *Frontiers in Education Conference*, 36th Annual. IEEE, p. 23-27.
- [29] C Kefalis, C Skordoulis, A Drigas 2025 Digital Simulations in STEM Education: Insights from Recent Empirical Studies, a Systematic Review *Encyclopedia MDPI* 5 (1), 10
- [30] M Karyotaki, A Drigas 2015 Online and other ICT applications for cognitive training and assessment *International Journal of Online Engineering (iJOE)* 11 (2), 36-42
- [31] Drigas AS, Argyri K, Vrettaros J (2009) Decade review (1999-2009): artificial intelligence techniques in student modeling. In: *World Summit on Knowledge Society*. Springer, pp 552–564
- [32] E Mitsea, N Lytra, A Akrivopoulou, A Drigas 2020 Metacognition, Mindfulness and Robots for Autism Inclusion. *Int. J. Recent Contributions Eng. Sci. IT* 8 (2), 4-20

- [33] Kokkalia, G., Drigas, A., & Economou, A. (2016). The role of games in special preschool education. *International Journal of Emerging Technologies in Learning (ijET)*, 11(12), 30-35.
- [34] A Drigas, C Papoutsi 2021 Nine Layer Pyramid Model Questionnaire for Emotional Intelligence. *Int. J. Online Biomed. Eng.* 17 (7), 123
- [35] Drigas, A. S., & Karyotaki, M. (2019). A Layered Model of Human Consciousness. *International Journal of Recent Contributions from Engineering, Science & IT (ijES)*, 7(3), 41- 50. <https://doi.org/10.3991/ijes.v7i3.11117>
- [36] A Drigas, M Karyotaki 2019 Attention and its Role: Theories and Models. *International Journal of Emerging Technologies in Learning* 14 (12), 169-182